

Published 1989

# Contributions from Breeding Forage and Turf Grasses



# Contributions from Breeding Forage and Turf Grasses

Proceedings of a symposium sponsored by Divisions C-1, C-5, and C-6 of  
the Crop Science Society of America in Atlanta, Georgia, 1 Dec. 1987.

## Editors

D. A. Sleper, K. H. Asay, and J. F. Pedersen

### *Organizing Committee*

D. A. Sleper

K. H. Asay

### *Editorial Committee*

D. A. Sleper

K. H. Asay

J. F. Pedersen

### *Editor-in-Chief CSSA*

C. W. Stuber

### *Managing Editor*

S. H. Mickelson

CSSA Special Publication Number 15

**Crop Science Society of America, Inc.**  
**Madison, Wisconsin, USA**  
**1989**

Cover Design: Patricia Jeffson

Crop Science Society of America, Inc.  
677 South Segoe Road, Madison, WI 53711, USA

Copyright © 1989 by the Crop Science Society of America, Inc.

**ALL RIGHTS RESERVED UNDER THE U.S. COPYRIGHT LAW  
OF 1978 (P.L. 94-553)**

Any and all uses beyond the limitations of the “fair use” provision of the law require written permission from the publisher(s) and/or the author(s); not applicable to contributions prepared by officers or employees of the U.S. Government as part of their official duties.

**Library of Congress Cataloging-in-Publication Data**

Printed in the United States of America.

# CONTENTS

Foreword .....	vii
Preface .....	ix
Contributors .....	xi
Conversion Factors for SI and non-SI units .....	xiii
1 Contributions from Breeding Forage and Turf Grasses—An Overview K. H. Asay and D. A. Sleper .....	1
2 Cool-Season Forage Grass Breeding: Progress, Potentials, and Benefits R. E. Barker and R. R. Kalton .....	5
3 Progress and Benefits to Humanity from Breeding Warm-Season Forage Grasses Glenn W. Burton .....	21
4 Progress and Benefits to Humanity from Breeding Cool-Season Grasses for Turf William A. Meyer and C. Reed Funk .....	31
5 Progress and Benefits to Humanity from Breeding Warm-Season Grasses for Turf Philip Busey .....	49
6 Breeding Forage Grasses to Maximize Animal Performance Gordon C. Marten .....	71
7 Breeding Grasses for the Future K. P. Vogel, H. J. Gorz, and F. A. Haskins .....	105



# FOREWORD

Among plants, the Gramineae, or grass family, provides the major component of human foods, either directly as cereal grains or indirectly by meat animals that graze the land or are provided grasses such as hay, silage, or freshly cut forage. Native stands of grasses throughout the world have sustained wildlife and later domesticated animals. Some of the wild grasses contributed genes to important cereal grains. Others have been collected and transported to gardens for evaluation and new uses. Still others have been used in hybridization for selection of new characteristics. Thus in less than 100 yr, the world's agriculture has benefitted from genetically improved grasses. Likewise, grasses have been selected for intensely managed and heavily used sites such as sports fields and parks.

The versatility of grasses is unsurpassed among plants. This small book provides documentation for some of the advances made in genetic improvement of grasses. The smallness of the book and the relatively few scientists who have dedicated their professional careers to grass breeding severely underplay the great importance of the contributions of grass breeding to society. It does, however, provide a benchmark upon which future gains may be judged. It also establishes the advances made by traditional plant breeding practices and emphasizes the genetic complexity in a cultivar adapted to special conditions. This complexity of adaptation must be recognized as molecular-based plant breeding emerges as a reality.

The organizers of the symposium are complimented for the timeliness of their efforts. It is hoped that this book will stimulate further investments in research on grass breeding because the opportunities are great.

Calvin O. Qualset, *president*  
Crop Science Society of America





## PREFACE

What are the contributions to humankind as the result of breeding forage and turf grasses? This publication attempts to answer this question as we look back over the history of grass breeding in the United States. The contributions from forage grass breeding have resulted in significant improvements in herbage and seed yield, insect and disease resistance, herbage quality, and animal performance, to mention a few. Turf grass breeding can also claim significant advancements in the areas of adaptation of species to environmental stresses, insect and disease resistance, higher seed yields, and better turf quality for a multitude of uses.

Forage and turf grass breeders have a rich history of cooperating with scientists from other disciplines to accomplish their research objectives. Within the chapters of this publication, it is apparent that significant contributions by the forage grass breeder have come about through cooperation with agronomists, animal scientists, pasture management specialists, economists, plant physiologists, entomologists, plant pathologists, and forage chemists. Likewise, the turf breeding effort has been enhanced by receiving cooperation from horticulturalists, irrigation specialists, plant pathologists, entomologists, and others. Collectively, their efforts have resulted in forage and turf grasses being established on vast acreages in the United States—far more than for any other individual crop species.

The breeding and genetics of forage and turf grasses has not received the amount of research support that many cash crops have in the United States. Vast acreages of forage and turf grasses are grown each year which make significant contributions to the livestock industry, wildlife conservation, soil and water conservation, recreation, and have tremendous aesthetic value, yet the amount of dollars spent and the number of scientists involved is far less than what is needed. With this impetus, the Grass Breeder's Work Planning Conference appointed a committee to organize a symposium to document the contributions made by the forage and turf grass breeders. The symposium was held at the annual meetings of the Crop Science Society of America (CSSA) at Atlanta, GA in 1987. This publication is the result of that symposium.

The most important attribute of any publication is the quality of its authorship. The authors of this publication are recognized as experts in their respective scientific disciplines. A great deal of gratitude is extended to the authors for their scholarly writings and enthusiasm for the topic at hand. Thanks is also due to the CSSA for sponsoring this important publication.

Editors:

D. A. Sleper  
K. H. Asay  
J. F. Pedersen



# CONTRIBUTORS

- K. H. Asay** Research Geneticist, USDA-ARS, Forage and Range Research Laboratory, Utah State University, Logan, Utah
- R. E. Barker** Research Geneticist, USDA-ARS, National Forage Seed Production Research Center, Corvallis, Oregon. Formerly with USDA-ARS, Northern Great Plains Research Laboratory, Mandan, North Dakota
- Glenn W. Burton** Research Geneticist, USDA-ARS and University of Georgia, Coastal Plain Experiment Station, Tifton, Georgia
- Philip Busey** Associate Professor of Ornamental Horticulture (Turfgrass Breeder), University of Florida, Fort Lauderdale, Florida
- C. Reed Funk** Professor, New Jersey Agricultural Experiment Station, Soils and Crops Department, Cook College, Rutgers University, New Brunswick, New Jersey
- H. J. Gorz** Supervisory Research Geneticist, USDA-ARS, Department of Agronomy, University of Nebraska, Lincoln, Nebraska
- F. A. Haskins** George Holmes Professor of Agronomy, Department of Agronomy, University of Nebraska, Lincoln, Nebraska
- R. R. Kalton** Forage Breeder, VISTA, Land O'Lakes Research Farm, Webster City, Iowa
- Gordon C. Marten** Research Agronomist and Professor, USDA-ARS and University of Minnesota, St. Paul, Minnesota
- William A. Meyer** President—Turfgrass Breeder, Pure Seed Testing, Inc., Hubbard, Oregon
- D. A. Sleper** Professor of Agronomy, Department of Agronomy, University of Missouri, Columbia, Missouri
- K. P. Vogel** Research Geneticist, USDA-ARS, Department of Agronomy, University of Nebraska, Lincoln, Nebraska



## **Conversion Factors for SI and non-SI Units**

## Conversion Factors for SI and non-SI Units

To convert Column 1 into Column 2, multiply by	Column 1 SI Unit	Column 2 non-SI Unit	To convert Column 2 into Column 1, multiply by
	<b>Length</b>		
0.621	kilometer, km ( $10^3$ m)	mile, mi	1.609
1.094	meter, m	yard, yd	0.914
3.28	meter, m	foot, ft	0.304
1.0	micrometer, $\mu\text{m}$ ( $10^{-6}$ m)	micron, $\mu$	1.0
$3.94 \times 10^{-2}$	millimeter, mm ( $10^{-3}$ m)	inch, in	25.4
10	nanometer, nm ( $10^{-9}$ m)	Angstrom, Å	0.1
	<b>Area</b>		
2.47	hectare, ha	acre	0.405
247	square kilometer, $\text{km}^2$ ( $10^3$ m) <sup>2</sup>	acre	$4.05 \times 10^{-3}$
0.386	square kilometer, $\text{km}^2$ ( $10^3$ m) <sup>2</sup>	square mile, mi <sup>2</sup>	2.590
$2.47 \times 10^{-4}$	square meter, m <sup>2</sup>	acre	$4.05 \times 10^3$
10.76	square meter, m <sup>2</sup>	square foot, ft <sup>2</sup>	$9.29 \times 10^{-2}$
$1.55 \times 10^{-3}$	square millimeter, $\text{mm}^2$ ( $10^{-6}$ m) <sup>2</sup>	square inch, in <sup>2</sup>	645
	<b>Volume</b>		
$9.73 \times 10^{-3}$	cubic meter, m <sup>3</sup>	acre-inch	102.8
35.3	cubic meter, m <sup>3</sup>	cubic foot, ft <sup>3</sup>	$2.83 \times 10^{-2}$
$6.10 \times 10^4$	cubic meter, m <sup>3</sup>	cubic inch, in <sup>3</sup>	$1.64 \times 10^{-5}$
$2.84 \times 10^{-2}$	liter, L ( $10^{-3}$ m <sup>3</sup> )	bushel, bu	35.24
1.057	liter, L ( $10^{-3}$ m <sup>3</sup> )	quart (liquid), qt	0.946
$3.53 \times 10^{-2}$	liter, L ( $10^{-3}$ m <sup>3</sup> )	cubic foot, ft <sup>3</sup>	28.3
0.265	liter, L ( $10^{-3}$ m <sup>3</sup> )	gallon	3.78
33.78	liter, L ( $10^{-3}$ m <sup>3</sup> )	ounce (fluid), oz	$2.96 \times 10^{-2}$
2.11	liter, L ( $10^{-3}$ m <sup>3</sup> )	pint (fluid), pt	0.473

Mass

2.20 × 10 <sup>-3</sup>	gram, g (10 <sup>-3</sup> kg)	454	ounce (avdp), oz
3.52 × 10 <sup>-2</sup>	gram, g (10 <sup>-3</sup> kg)	28.4	ounce (avdp), oz
2.205	kilogram, kg	0.454	pound, lb
0.01	kilogram, kg	100	quintal (metric), q
1.10 × 10 <sup>-3</sup>	kilogram, kg	907	ton (2000 lb), ton
1.102	megagram, Mg (tonne)	0.907	ton (U.S.), ton
1.102	tonne, t	0.907	ton (U.S.), ton

Yield and Rate

0.893	kilogram per hectare, kg ha <sup>-1</sup>	1.12	pound per acre, lb acre <sup>-1</sup>
7.77 × 10 <sup>-2</sup>	kilogram per cubic meter, kg m <sup>-3</sup>	12.87	pound per bushel, bu <sup>-1</sup>
1.49 × 10 <sup>-2</sup>	kilogram per hectare, kg ha <sup>-1</sup>	67.19	bushel per acre, 60 lb
1.59 × 10 <sup>-2</sup>	kilogram per hectare, kg ha <sup>-1</sup>	62.71	bushel per acre, 56 lb
1.86 × 10 <sup>-2</sup>	kilogram per hectare, kg ha <sup>-1</sup>	53.75	bushel per acre, 48 lb
0.107	liter per hectare, L ha <sup>-1</sup>	9.35	gallon per acre
893	tonnes per hectare, t ha <sup>-1</sup>	1.12 × 10 <sup>-3</sup>	pound per acre, lb acre <sup>-1</sup>
893	megagram per hectare, Mg ha <sup>-1</sup>	1.12 × 10 <sup>-3</sup>	pound per acre, lb acre <sup>-1</sup>
0.446	megagram per hectare, Mg ha <sup>-1</sup>	2.24	ton (2000 lb) per acre, ton acre <sup>-1</sup>
2.24	meter per second, m s <sup>-1</sup>	0.447	mile per hour

Specific Surface

10	square meter per kilogram, m <sup>2</sup> kg <sup>-1</sup>	0.1	square centimeter per gram, cm <sup>2</sup> g <sup>-1</sup>
1000	square meter per kilogram, m <sup>2</sup> kg <sup>-1</sup>	0.001	square millimeter per gram, mm <sup>2</sup> g <sup>-1</sup>

Pressure

9.90	megapascal, MPa (10 <sup>6</sup> Pa)	0.101	atmosphere
10	megapascal, MPa (10 <sup>6</sup> Pa)	0.1	bar
1.00	megagram per cubic meter, Mg m <sup>-3</sup>	1.00	gram per cubic centimeter, g cm <sup>-3</sup>
2.09 × 10 <sup>-2</sup>	pascal, Pa	47.9	pound per square foot, lb ft <sup>-2</sup>
1.45 × 10 <sup>-4</sup>	pascal, Pa	6.90 × 10 <sup>3</sup>	pound per square inch, lb in <sup>-2</sup>

continued on next page

## Conversion Factors for SI and non-SI Units

To convert Column 1 into Column 2, multiply by	Column 1 SI Unit	Column 2 non-SI Unit	To convert Column 2 into Column 1, multiply by
	<b>Temperature</b>		
	Kelvin, K	Celsius, °C	Celsius, °C
	Celsius, °C	Fahrenheit, °F	Fahrenheit, °F
1.00 (K - 273) (9/5 °C) + 32			1.00 (°C + 273) 5/9 (°F - 32)
	<b>Energy, Work, Quantity of Heat</b>		
	joule, J	British thermal unit, Btu	1.05 × 10 <sup>3</sup>
9.52 × 10 <sup>-4</sup>		calorie, cal	4.19
0.239		erg	10 <sup>-7</sup>
10 <sup>7</sup>		foot-pound	1.36
0.735		calorie per square centimeter (langley)	4.19 × 10 <sup>4</sup>
2.387 × 10 <sup>-5</sup>		dyne	10 <sup>-5</sup>
10 <sup>5</sup>		calorie per square centimeter minute (irradiance), cal cm <sup>-2</sup> min <sup>-1</sup>	698
1.43 × 10 <sup>-3</sup>			
	<b>Transpiration and Photosynthesis</b>		
	milligram per square meter second, mg m <sup>-2</sup> s <sup>-1</sup>	gram per square decimeter hour, g dm <sup>-2</sup> h <sup>-1</sup>	27.8
3.60 × 10 <sup>-2</sup>		micromole (H <sub>2</sub> O) per square centi- meter second, μmol cm <sup>-2</sup> s <sup>-1</sup>	180
5.56 × 10 <sup>-3</sup>		milligram per square centimeter second, mg cm <sup>-2</sup> s <sup>-1</sup>	10 <sup>4</sup>
10 <sup>-4</sup>		milligram per square decimeter hour, mg dm <sup>-2</sup> h <sup>-1</sup>	2.78 × 10 <sup>-2</sup>
35.97			
	<b>Plane Angle</b>		
	radian, rad	degrees (angle), °	1.75 × 10 <sup>-2</sup>
57.3			



Electrical Conductivity, Electricity, and Magnetism

10	siemen per meter, S m <sup>-1</sup>	millimho per centimeter, mmho cm <sup>-1</sup>	0.1
10 <sup>4</sup>	tesla, T	gauss, G	10 <sup>-4</sup>
<b>Water Measurement</b>			
9.73 × 10 <sup>-3</sup>	cubic meter, m <sup>3</sup>	acre-inches, acre-in	102.8
9.81 × 10 <sup>-3</sup>	cubic meter per hour, m <sup>3</sup> h <sup>-1</sup>	cubic feet per second, ft <sup>3</sup> s <sup>-1</sup>	101.9
4.40	cubic meter per hour, m <sup>3</sup> h <sup>-1</sup>	U.S. gallons per minute, gal min <sup>-1</sup>	0.227
8.11	hectare-meters, ha-m	acre-feet, acre-ft	0.123
97.28	hectare-meters, ha-m	acre-inches, acre-in	1.03 × 10 <sup>-2</sup>
8.1 × 10 <sup>-2</sup>	hectare-centimeters, ha-cm	acre-feet, acre-ft	12.33

Concentrations

1	centimole per kilogram, cmol kg <sup>-1</sup>	milliequivalents per 100 grams, meq	1
	(ion exchange capacity)	100 g <sup>-1</sup>	
0.1	gram per kilogram, g kg <sup>-1</sup>	percent, %	10
1	milligram per kilogram, mg kg <sup>-1</sup>	parts per million, ppm	1

Radioactivity

2.7 × 10 <sup>-11</sup>	becquerel, Bq	curie, Ci	3.7 × 10 <sup>10</sup>
2.7 × 10 <sup>-2</sup>	becquerel per kilogram, Bq kg <sup>-1</sup>	picocurie per gram, pCi g <sup>-1</sup>	37
100	gray, Gy (absorbed dose)	rad, rd	0.01
100	sievert, Sv (equivalent dose)	rem (roentgen equivalent man)	0.01

Plant Nutrient Conversion

<i>Elemental</i>		<i>Oxide</i>	
2.29	P	P <sub>2</sub> O <sub>5</sub>	0.437
1.20	K	K <sub>2</sub> O	0.830
1.39	Ca	CaO	0.715
1.66	Mg	MgO	0.602