

Fast-Track Introgression of “QTL-hotspot” for Root Traits and Other Drought Tolerance Traits in JG 11, an Elite and Leading Variety of Chickpea

The Plant Genome 6, doi: 10.3835/plantgenome2013.07.0022

A calculation error for standard deviation was made in the original version of this manuscript. Corrections were made in three instances as follows. ■

Abstract: Root trait phenotyping of these IIs showed higher rooting depth (RDp) (average 115.21 ± 2.24 cm) in all 29 IIs, better root length density (RLD) (average 0.41 ± 0.02 cm cm⁻³) in 26 IIs, and higher root dry weight (RDW) (average 1.25 ± 0.08 g per cylinder) as compared to the recurrent parent, JG 11 (111.70 cm for RDp, 0.39 cm cm⁻³ for RLD, and 1.10 g per cylinder for RDW), as well as the donor parent, ICC 4958 (114.20 cm for RDp, 0.45 cm cm⁻³ for RLD, and 1.25 g per cylinder for RDW).

Page 5: In fact, 26 out of 29 IIs showed better RLD as an average 0.41 ± 0.02 cm cm⁻³ in comparison to the recurrent parent JG 11 (0.39 cm cm⁻³).

Last line of Table 4:

SD±	6.43±1.19	2.24±0.40	0.02±0.00	0.08±0.01
-----	-----------	-----------	-----------	-----------

Published in The Plant Genome 9.
doi: 10.3835/plantgenome2013.07.0022er

© Crop Science Society of America
5585 Guilford Rd., Madison, WI 53711 USA
This is an open access article distributed under the CC BY-NC-ND
license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).